

Amplitude analysis of GlueX phase 1 $(p\eta'\pi^0)$ data using waveset from VMD (Vector Meson Dominance) model for a_2 photoproduction

Florida International University 2021

Mariana Khachatryan

1. Fitting entire GlueX phase 1 data (adding fall 2018) for four γ polarization plane angles relative to horizontal ($0, 45, 90, 135^\circ$) + amorphous data using loop statement in AmpTools
2. Fit intensity with a waveset that includes S, D waves, where we include only certain M and ϵ values for D waves that are expected to have nonzero contribution based on Vincents VMD model (Vector Meson Dominance):

- $S_0^{(+)}, S_0^{(-)}, D_0^{(+)}, D_1^{(+)}, D_2^{(+)}, D_{-1}^{(-)}, D_0^{(-)}, D_1^{(-)}$

$$s = m_N^2 + 2m_N E_\gamma, E_\gamma = 8.5 \text{ GeV and } t = -0.2 \text{ GeV}^2,$$

$${}^{(+)}[D]_{m=2} = 2.35$$

$${}^{(+)}[D]_{m=1} = 0.82$$

$${}^{(+)}[D]_{m=0} = 0.11$$

$${}^{(-)}[D]_{m=1} = 0.84$$

$${}^{(-)}[D]_{m=0} = 0.45$$

$${}^{(-)}[D]_{m=-1} = 0.09$$



Photoproduction of $a_2(1320) \rightarrow \eta\pi^0$
 Vincent Mathieu
 May 26, 2021

3. Invariant mass bin size of 75 MeV/c², momentum transfer bin size of 0.6 (GeV/c)²

4. Hopefully, the increased statistics and decreased number of fit parameters will allow to also calculate uncertainties from bootstrapping method

Plot acceptance uncorrected results


Acceptance uncorrected




Fit with $S_0, P_{0,1}, D_{0,1,2}$ $\varepsilon = \pm 1$

Acceptance uncorrected


D0pl




D1p




D2pl




Dmi1mi



D0mi




D1mi







All waves





Total



$0.1 < t < 0.7 \text{ (GeV/c)}^2$


Uncertainties from bootstrapping



0.1< t < 0.7 (GeV/c)² Uncertainties from bootstrapping


$0.1 < t < 0.7 \text{ (GeV/c}^2)$

Uncertainties from bootstrapping




$1.1 \quad 1.3 \quad 1.5 \quad 1.7 \quad 1.9$

$M(\eta'\pi^0)$ [GeV/c²]




$1.1 \quad 1.3 \quad 1.5 \quad 1.7 \quad 1.9$

$M(\eta'\pi^0)$ [GeV/c²]



$1.1 \quad 1.3 \quad 1.5 \quad 1.7 \quad 1.9$

$M(\eta'\pi^0)$ [GeV/c²]



$1.1 \quad 1.3 \quad 1.5 \quad 1.7 \quad 1.9$

$M(\eta'\pi^0)$ [GeV/c²]

Moments calculated using fitted partial waves

Uncertainties from bootstrapping

$0.1 < t < 0.7 \text{ (GeV/c)}^2$

